

Direct Wavelet Expansion of the Primordial Power Spectrum

Pia Mukherjee¹, and Yun Wang¹

ABSTRACT

In order to constrain and possibly detect unusual physics during inflation, we allow the power spectrum of primordial matter density fluctuations, $P_{in}(k)$, to be an arbitrary function in the estimation of cosmological parameters from data. The multi-resolution and good localization properties of orthogonal wavelets make them suitable for detecting features in $P_{in}(k)$. We expand $P_{in}(k)$ directly in wavelet basis functions. The likelihood of the data is thus a function of the wavelet coefficients of $P_{in}(k)$, as well as the Hubble constant H_0 , baryon density $\Omega_b h^2$, cold dark matter density $\Omega_c h^2$ and the reionization optical depth τ_{ri} , in a flat Λ CDM cosmology. We derive constraints on these parameters from Cosmic Microwave Background (CMB) anisotropy data (WMAP, CBI, and ACBAR) and large scale structure (LSS) data (2dFGRS and PSCZ) using the Markov Chain Monte Carlo (MCMC) technique. The direct wavelet expansion method is different and complimentary to the wavelet band power method of Mukherjee & Wang (2003a,b), and results from the two methods are consistent. In addition, as we demonstrate, the direct wavelet expansion method has the advantage that once the wavelet coefficients have been constrained, the reconstruction of $P_{in}(k)$ can be effectively denoised, i.e., $P_{in}(k)$ can be reconstructed using only the coefficients that, say, deviate from zero at greater than 1σ . In doing so the essential properties of $P_{in}(k)$ are retained. The reconstruction also suffers much less from the correlated errors of binning methods. The shape of the primordial power spectrum, as reconstructed in detail here, reveals an interesting new feature at $0.001 \lesssim k/\text{Mpc}^{-1} \lesssim 0.005$. It will be interesting to see if this feature is confirmed by future data. The reconstructed and denoised $P_{in}(k)$ is favored over the scale-invariant and power-law forms at $\gtrsim 1\sigma$.

1. Introduction

In spite of the success of the inflationary paradigm (Guth 1981; Albrecht & Steinhardt 1982; Gott 1982; Linde 1983; Kolb & Turner 1990; Hu 2003; Peebles & Ratra 2003), we are

¹Department of Physics & Astronomy, Univ. of Oklahoma, 440 W Brooks St., Norman, OK 73019; email: pia,wang@nhn.ou.edu

still in search of the correct model of inflation. The simplest models of inflation predict a power-law primordial matter power spectrum (for example, Linde (1983); Freese, Frieman, & Olinto (1990); La & Steinhardt (1991)). However, there are many other viable models of inflation which predict primordial power spectra which cannot be parametrized by a simple power law (for example, Holman et al. (1991ab); Wang (1994); Randall, Soljagic, & Guth (1996); Adams, Ross, & Sarkar (1997); Lesgourgues, Polarski, & Starobinsky (1997)). These can represent unusual physics in the very early universe (for example, see Chung et al. (2000); Enqvist & Kurki-Suonio (2000); Lyth, Ungarelli, & Wands (2002); Feng & Zhang (2003)), and then assuming a power-law primordial matter power spectrum could lead to our missing the discovery of the possible features in it and erroneous estimates of the cosmological parameters (Kinney 2001).

With results from WMAP, CMB data continue to be consistent with inflation (Bennett et al. 2003, Spergel et al. 2003), having even detected the predicted anti-correlation between CMB temperature and polarization fluctuations near l of 150. While the derived cosmological parameter constraints are consistent with previous constraints from independent and complementary observations, Peiris et al. (2003) report that the data, both with and without the use of complementary LSS data, seem to indicate a preferred scale in the primordial power spectrum. They conclude that the data suggest at the 2σ level, but do not require, that the scalar spectral index runs from $n_s > 1$ on large scales to $n_s < 1$ on small scales. Their analysis was done assuming a primordial power spectrum of power-law form about a pivot scale of 0.05 Mpc^{-1} (0.002 Mpc^{-1}) when using scalar (both scalar and tensor) modes.

Now that such precision measurements of CMB anisotropy have been made, we aim to allow the primordial power spectrum to be an arbitrary function, independent of any particular inflationary model, and use the data to place constraints on its shape. WMAP data underscore the importance of such measurements suggested earlier by Wang, Spergel, & Strauss (1999); Wang & Mathews (2002) who used conventional top-hat binning and linear interpolation respectively in the parametrization of $P_{in}(k)$. Mukherjee & Wang (2003a) parametrized $P_{in}(k)$ in terms of wavelet band powers, and arrived at a detailed measurement of $P_{in}(k)$ using pre-WMAP CMB data. A preliminary indication of a feature was reported. The method was shown to work better than the linear-interpolation method of Wang & Mathews (2002). Mukherjee & Wang (2003b) (hereafter MW03b) present constraints on $P_{in}(k)$ from WMAP data using the wavelet band powers method, and also the tophat binning method for comparison. It was found that the reconstructed spectrum deviates from scale-invariant as well as power-law models by approximately 1σ . While the data do not rule out simple parametrizations such as scale-invariance, as also concluded by Bridle et al. (2003), Barger, Lee & Marfatia (2003), Seljak, McDonald & Makarov (2003), there is indication of

excess power or a preferred scale at $k \sim 0.01 \text{ Mpc}^{-1}$ (Peiris et al. 2003, Spergel et al. 2003, MW03b). Wang & Mathews (2002); Mukherjee & Wang (2003a) show that pre-MAP data indicated a similar feature.

It is worth investigating this issue further. Since data in CMB multipole space is essentially being used to constrain $P_{in}(k)$, its band powers as estimated from the data become correlated with one another (because of the cosmological model dependent non-linear mapping between wavenumber k and multipole l , see Tegmark & Zaldarriaga 2002). The $P_{in}(k)$ parameters in any model parametrization are correlated with the cosmological model parameters (that describe the evolution of structure from the initial conditions to the present). These correlations can make it tricky to efficiently constrain $P_{in}(k)$.

In this paper, we explore a more direct method of using wavelets to probe $P_{in}(k)$; we parametrize $P_{in}(k)$ by the coefficients of its wavelet expansion. We obtain constraints on these wavelet coefficients and cosmological parameters from current CMB and LSS data. This direct wavelet expansion method is different and complimentary to the wavelet band power method of MW03a,b.² It has the advantage that once the wavelet coefficients have been constrained, the reconstruction of $P_{in}(k)$ can be effectively denoised, i.e., $P_{in}(k)$ can be reconstructed using only the coefficients that deviate from zero at greater than 1σ . In doing so the essential properties of $P_{in}(k)$ are retained. Because wavelets are adaptive, and the wavelet basis complete, any large scale as well as localized features in $P_{in}(k)$ can be efficiently picked up by a small number of coefficients. Such properties make wavelets useful tools in signal reconstruction. This method is thus used to reconstruct $P_{in}(k)$ as an arbitrary function over the entire range in k that cosmological data are sensitive to, and since it can be reliably reconstructed from a small number of wavelet coefficients, it suffers much less from the correlated errors of binning methods.

We describe our method in Sec.2. We present our results in Sec.3. Sec.4 contains a summary and discussion.

²The direct wavelet expansion method uses a set of orthogonal basis functions that are very different from those used in binning methods; the resultant constraints on $P_{in}(k)$ can be thus checked for consistency. This can be significant because parameter correlations, which are important for this problem, are very different for the different sets of basis functions. We found the parameter correlations to be the smallest in the direct wavelet expansion method.

2. Method

We expand the primordial power spectrum $P_{in}(k)$ ³ directly in wavelet basis functions,

$$P_{in}(k_i) = \sum_{j=0}^{J-1} \sum_{l=0}^{2^j-1} b_{j,l} \psi_{j,l}(k_i), \quad (1)$$

where $\psi_{j,l}$ are the wavelet basis functions, constructed from the dilations and translations of a mother function $\psi(k)$, via (Daubechies 1992; Press et al. 1994)

$$\psi_{j,l}(k) = \left(\frac{2^j}{L}\right)^{1/2} \psi(2^j k/L - l). \quad (2)$$

The resulting wavelet bases are discrete, compactly supported and orthogonal with respect to both the scale j and the position l indices, and hence their coefficients provide a complete and non-redundant (hence invertible) representation of the function $P_{in}(k)$. The scale index j increases from 0 to $J - 1$, and wavelet coefficients with increasing j represent structure in $P_{in}(k)$ on increasingly smaller scales, with each scale a factor of 2 finer than the previous one. The index l , which runs from 0 to $2^j - 1$ for each j , denotes the position of the wavelet basis $\psi_{j,l}$ within the j th scale. (Note that the properties of the basis functions are completely different from the basis functions in binning methods). Therefore the wavelet coefficients in Eq.(1), together with the scaling function coefficient which represents the mean of the function, (denoted $a_{0,0}$), total to 2^J in number; correspondingly the function $P_{in}(k)$ is reconstructed at 2^J points. We have considered the case when these sample points k_i , over which the wavelet basis is defined and at which the function $P_{in}(k)$ is reconstructed, are equally spaced in $\log(k)$. We use 16 wavelet coefficients to represent $P_{in}(k)$ in the range $0.0002 \lesssim k/(\text{Mpc}^{-1}) \lesssim 0.2$ that the data are sensitive to, with the spectrum for $k < 0.0002 \text{ Mpc}^{-1}$ set to $P_{in}(k = 0.0002 \text{ Mpc}^{-1})$ and the spectrum for $k > 0.2 \text{ Mpc}^{-1}$ set to $P_{in}(k = 0.2 \text{ Mpc}^{-1})$.

Thus, when the sample points are equally spaced, the coefficient $b_{j,l}$ measures the signal in $P_{in}(k)$ on scale $L/2^j$, and centered at $k = lL/2^j$, where L is the full length over which $P_{in}(k)$ is being expanded. The simultaneous and adaptive localization of the wavelet basis functions in position and frequency space make them useful tools for analyzing a variety of data. The information content of most signals can be efficiently encoded in wavelet space; hence the wavelet representation of a signal is often sparse and allows for significant compression (as coefficients below a threshold can be thrown away with very little loss of information making

³For a power-law primordial power spectrum, $P_{\mathcal{R}} = 2.95 \times 10^{-9} A (k/k_0)^{n_s-1}$ (Spergel et al. 2003). For an arbitrary primordial power spectrum, we define $P_{in}(k) = \frac{P_{\mathcal{R}}}{2.95 \times 10^{-9}}$.

them important tools in denoising). For example a scale-invariant $P_{in}(k)$ requires only two non-zero coefficients (the scaling function coefficient, and the wavelet coefficient of scale 0), and when attempting to reconstruct this function from noisy data all other wavelet coefficients would be zero within error bars. If some other wavelet coefficients are non-zero then that implies additional structure in the function.

Thus our goal is to constrain $P_{in}(k)$ by estimating its wavelet coefficients, together with cosmological parameters, from a likelihood analysis of cosmological data. We do not impose any particular inflation model dependent parametrization on $P_{in}(k)$, and allow it to be an arbitrary function. After constraining the wavelet coefficients, marginalizing over all other parameters considered, we can choose to reconstruct $P_{in}(k)$ using only those wavelet coefficients that are significantly constrained. In doing so all the important features of $P_{in}(k)$ are retained and one is essentially denoising. Thus in this construction features in any $P_{in}(k)$ can be localized in scale j and position l with the help of a few coefficients, and the resulting reconstruction also suffers much less from the correlated errors of binning methods.

This direct wavelet expansion method is very different in nature from the wavelet band power method used in MW03a. In the wavelet band power method of MW03a, the primordial power spectrum is parametrized in terms of wavelet band powers; the window functions relating the wavelet band powers to the primordial power spectrum are the squares of the Fourier transforms of the wavelet basis functions (apart from a j dependent normalization factor). Wavelets that are more smooth and hence less localized in pixel space and correspondingly more localized in Fourier space, such as the Daubechies 20 wavelet are the suitable wavelets to use for this parametrization. In the direct wavelet expansion method we use the coefficients of a wavelet expansion of $P_{in}(k)$ to characterize it; the window functions in this case are the mutually orthogonal wavelet basis functions $\psi_{j,l}$. Thus here we need to use wavelets that are most localized in k space, which is the pixel space here, such as the Haar and the Daubechies 4 (hereafter D4) wavelet. Among such wavelets different wavelets give very similar results. Other differences between the two methods are that in the wavelet band power method the bin locations and their separations are not arbitrary, as explained further in MW03a (see also footnote 7 here), whereas here we can probe the function down to arbitrary resolution by reconstructing more coefficients. The correlations among the reconstructed parameters are different and smaller in the direct expansion method as compared to those in binning methods, and the reconstruction here can be effectively denoised; these properties have been elaborated on earlier in this section.

We also make use of the mutual independence of the wavelet basis functions in computing the projections of the CMB angular power spectrum C_l 's and the matter power spectrum $P(k)$ for each wavelet basis, thus avoiding calling CMBFAST (Seljak & Zaldarriaga 1996)

or CAMB (Lewis, Challinor, & Lasenby 2000), when only the wavelet coefficients are varied for the same cosmological parameters. This leads to a significant computational speed up. We write ⁴

$$\begin{aligned}
C_l(\{b_{j,l}\}, \mathbf{s}) &= (4\pi)^2 \int \frac{dk}{k} P_{in}(k) |\Delta_{Tl}(k, \tau = \tau_0)|^2 \\
&= \sum_{j,l} b_{j,l} \int \frac{dk}{k} \psi_{j,l}(k) |\Delta_{Tl}(k, \tau = \tau_0)|^2 \\
&\equiv \sum_{j,l} b_{j,l} C_l^{j,l}(\mathbf{s}),
\end{aligned} \tag{3}$$

where the cosmological model dependent transfer function $\Delta_{Tl}(k, \tau = \tau_0)$ is an integral over conformal time τ of the sources which generate CMB temperature fluctuations, τ_0 being the conformal time today, and $\mathbf{s}$ represents cosmological parameters other than the $b_{j,l}$'s. The $C_l^{j,l}(\mathbf{s})$'s are the wavelet projections of the C_l 's, computed here using CAMB⁵.

Similarly the wavelet projections of the evolved matter power spectrum $P(k)$ are computed for LSS data as follows,

$$\begin{aligned}
P(k, \{b_{j,l}\}, \mathbf{s}) &= P_{in}(k) * T(k)^2 \\
&= \sum_{j,l} b_{j,l} \psi_{j,l}(k) * T(k)^2 \\
&\equiv \sum_{j,l} b_{j,l} P^{j,l}(k, \mathbf{s})
\end{aligned} \tag{4}$$

where $T(k)$ is the matter transfer function.

Since the theoretical predictions to be compared with CMB and LSS data both depend on $P_{in}(k)$, and the cosmological parameters, they can together be used to place robust constraints on cosmological models and their initial conditions. Note however that as pointed out by Elgaroy, Gramann, & Lahav (2002), it is hard to detect features in the primordial power spectrum at $k < 0.03h\text{Mpc}^{-1}$ using LSS data only.

⁴In computing these wavelet projections, we note that $\psi_{k,l}$ are defined only at a finite number of points (here 16) and its value at any intermediate point is obtained by linear interpolation. This is appropriate; since we are using a discrete transform and correspondingly estimating a function at a finite number of points, no additional information at smaller scales is either assumed or included. We have checked that no interpolation errors are introduced at this stage.

⁵<http://camb.info/>

We use the Markov Chain Monte Carlo (MCMC) technique, illustrated for example in Lewis & Bridle (2002); Verde et al. (2003) to estimate the likelihood functions of the parameters. This technique is necessitated by the large number of parameters being varied here, and at the same time is free of the interpolation errors expected in the much slower grid based methods. At its best, the MCMC method scales approximately linearly with the number of parameters. See Neil (1993) for a review, and Hannestad (2000); Knox et al. (2001); Rubino-Martin et al. (2002) for other applications of this method to CMB data analysis. An MCMC sampler based on the Metropolis-Hastings algorithm has been made available in the software package CosmoMC (Lewis & Bridle 2002).

The MCMC method is used to trace the full posterior distribution of the parameters in a flat Λ CDM cosmology assuming wide uniform priors of $0.4 < h < 1.0$, where $h = H_0/(100 \text{ kms}^{-1} \text{ Mpc}^{-1})$, $0.005 < \Omega_b h^2 < 0.1$ and $0.1 < \Omega_c h^2 < 0.99$ on the cosmological parameters. The two wavelet coefficients that set the level of $P_{in}(k)$ are allowed to vary between 0 and 5, while the remaining wavelet coefficients are allowed to vary between ± 2.5 , with the requirement that the resulting $P_{in}(k)$ be positive definite⁶. We use a weak prior on the age of the universe of $t_0 > 10$ Gyrs. We have run several chains, and have checked for convergence and mixing. From the MCMC samples thus obtained, the marginalized posterior distributions and confidence limits of the parameters are estimated.

3. Results

We use CMB temperature anisotropy data from WMAP (Bennett et al. 2003), complemented at $l > 800$ by data from CBI (Pearson et al. 2002) and ACBAR (Kuo et al. 2002) upto an l_{max} of 2000, marginalizing analytically over known beamwidth and calibration uncertainties (Bridle et al. 2002). We use LSS data from 2dFGRS (Percival et al. 2002; the redshift space power spectrum is assumed to be proportional to the real space power spectrum on the large scales used) and PSCZ (Hamilton & Tegmark 2002; decorrelated real space power spectrum) galaxy redshift surveys over linear scales. A linear bias is assumed and analytically marginalized over (Bridle et al. 2002). LSS data are thus effectively being used to constrain the shape and not directly the amplitude of the matter power spectrum.

We assume Gaussian adiabatic scalar perturbations in a flat universe with a cosmological constant. We do not use tensor modes in this paper, since current data are not sensitive to tensor contributions. With freedom allowed in the shape of the primordial power spectrum,

⁶Note that for a scale-invariant $P_{in}(k)$ of unity the two coefficients that set the level are each 2.828 while the remaining 14 coefficients are 0.

the temperature anisotropy data themselves cannot distinguish well between different values of the reionization optical depth τ_{ri} . Hence we simply make use of the WMAP constraint of $\tau_{ri} = 0.17 \pm 0.04$, derived for a flat Λ CDM cosmology from WMAP’s TE polarization data (Kogut et al. 2003), where the error estimate includes systematic and foreground uncertainties. We marginalize the distributions of the other parameters over a Gaussian distribution in τ_{ri} centered at 0.17 with a standard deviation of 0.04. For this, we calculate the likelihood distributions of the wavelet coefficients as well as the Hubble constant H_0 , baryon density $\Omega_b h^2$ and the cold dark matter density $\Omega_c h^2$ at 5 select values of τ_{ri} and approximate the marginalization over τ_{ri} using the 5-point Gauss-Hermite quadrature summation.

Fig.1 shows the marginalized 1d distributions of the parameters, obtained from an MCMC based likelihood analysis of CMB and LSS data, after marginalizing over the assumed distribution for τ_{ri} . All the parameters are well constrained, lying well within the imposed priors.

Fig.2 shows the reconstructed $P_{in}(k)$ (solid curves) and its 1σ confidence region (dotted curves), obtained from the mean values and standard deviations of the wavelet coefficients. The uncertainty in $P_{in}(k)$ has been calculated using:

$$\sigma_{P_{in}(k_i)}^2 = \frac{1}{N} \sum \left[\sum_{j,l} (b_{j,l} - \bar{b}_{j,l}) \frac{\partial P_{in}(k_i)}{\partial b_{j,l}} \right]^2, \quad (5)$$

where the average is over the MCMC samples, $\bar{b}_{j,l}$ denotes the mean value of the wavelet coefficient, and $\frac{\partial P_{in}(k_i)}{\partial b_{j,l}} = \psi_{j,l}(k_i)$; correlations between the wavelet coefficients have thus been accounted for in computing the uncertainty in $P_{in}(k)$. The upper panel shows the reconstructed $P_{in}(k)$ using all wavelet coefficients. These constraints are shown overlaid on the constraints obtained by MW03b using the wavelet band power method.⁷ We see that while the constraints from the two methods are completely consistent, a new feature shows up at large angular scales around $0.001 \lesssim k/\text{Mpc}^{-1} \lesssim 0.005$. The lower panel shows the spectrum reconstructed using only those coefficients that deviate from 0 at $> 1\sigma$. These constraints are shown overlaid on the constraints obtained using the tophat binning method with bands carefully positioned closer together so as to be able to pick up the feature at large angular scales around $0.001 \lesssim k/\text{Mpc}^{-1} \lesssim 0.005$, and placed further apart in the remaining parts of the spectrum. The feature shows up consistently.

⁷Note that in the wavelet banding method the locations and the widths of the bands are not arbitrary. As explained in Mukherjee & Wang 2003a, the band power window functions in this case are the modulus squared of the Fourier transforms of the wavelet basis functions. By definition then, since each scale is a factor of 2 finer than the previous one, the bands lie $\log_{10}2$ apart in k .

The coefficients that are significant at the greater than 1σ level are $b_{0,0}$, $b_{1,0}$, $b_{2,0}$, $b_{2,1}$ and $b_{3,2}$, together with the scaling function coefficient. Notice that with 10 of the 16 wavelet coefficients dropped, the spectrum reconstructed in panel (b) still has all the essential features of the spectrum shown in panel (a). The rise in power that shows up near $k \sim 0.01 \text{ Mpc}^{-1}$ mimics a power-law model with possible running and is what was reported in MW03b and by WMAP. In this paper we report a localized new feature at $0.001 \lesssim k/\text{Mpc}^{-1} \lesssim 0.005$ that would be missed by a power-law parametrization, and by other model-dependent parametrizations. (We note that section 5 of Peiris et al. (2003) seems to indicate the same large scale feature obtained here, but they used the model dependent parametrization of Adams, Cresswell, & Easterher (2001) and constrained the $P_{in}(k)$ parameters at a fixed set of cosmological parameters.) This new feature would also be missed by a model-independent binning method if the bin separation were not small enough at the appropriate locations.⁸

In the direct expansion method the constraints obtained simultaneously on the cosmological parameters are $h = 0.59 \pm 0.09$, $\Omega_b h^2 = 0.018 \pm 0.004$ and $\Omega_c h^2 = 0.143 \pm 0.019$. In all the constraints presented here we have used the parameter mean values together with marginalized 1σ standard deviations.⁹

The χ^2_{eff} of the best fit model in the direct wavelet expansion method is 1032.45 for a total of 964 data points, when constraining 16 wavelet coefficients and 3 cosmological parameters with an external prior on τ_{ri} being imposed and marginalized over.¹⁰ A scale-invariant model with 5 constrained parameters has a χ^2_{eff} of 1044.33 for the same data. One can see that this model is equivalent to holding fixed the 14 wavelet coefficients that are responsible for any deviations from scale-invariance. Thus the difference in χ^2 between the two models has a χ^2 -distribution with 14 degrees of freedom. By computing the probability of obtaining such a χ^2 we are able to compare the two models and find that the scale-invariant model is disfavoured at $\sim 0.6\sigma$. The χ^2_{eff} of the best fit model in the direct wavelet expansion method using only the 6 significantly constrained wavelet coefficients is 1039.52 for 9 constrained parameters. This is favored over the scale-invariant model at

⁸Note that a uniform binning with a small bin size can lead to oversampling of the data, which would result in large correlated error bars for the estimated bin amplitudes, thus burying any possible features (Bridle et al. 2003).

⁹As discussed in Lewis & Bridle (2002), the MCMC samples from the posterior do not provide accurate estimates of parameter best-fit values, because in higher dimensions the best-fit region typically has a much higher likelihood than the mean but occupies a very small fraction of parameter space.

¹⁰The number of data points in the different data sets used are 899 (WMAP), 4 (CBI), 7 (ACBAR), 32 (2dFGRS) and 22 (PSCZ).

$\gtrsim 1\sigma$.¹¹ Note that in our analysis the number of constrained parameters are large and the significance of deviations from scale-invariance are small because we allow $P_{in}(k)$ to be an arbitrary function over the entire k range. While the deviation in the shape of $P_{in}(k)$ from scale-invariance is not found to be significant at this stage, the primordial power spectrum reconstructed here seems to indicate intriguing features that could indicate either unusual inflationary physics or hidden systematics in current CMB data.

4. Conclusions

It is important to reconstruct the primordial power spectrum as a free function from cosmological data, the aim being to test the assumptions that are usually made about $P_{in}(k)$, and to constrain it model-independently so as to be able to detect any features in it that may be signatures of unusual physics during inflation. We have presented a new and powerful method to extract the primordial power spectrum $P_{in}(k)$ as an arbitrary function from observational data — the direct expansion of $P_{in}(k)$ in wavelet basis functions.

The wavelet basis functions afford an efficient (sparse) and non-redundant (invertible) expansion of a function, with coefficients that encode information about the structure of the function on different scales, and with adaptive position resolution. These properties make wavelets suitable tools for directly probing the primordial power spectrum for features. Reconstructing $P_{in}(k)$ using only the wavelet coefficients that deviate from 0 at $> 1\sigma$ amounts to denoising the resultant power spectrum while retaining all its important features (Fig 2). The direct wavelet expansion method also yields smaller parameter correlations than the binning methods.

The $P_{in}(k)$ reconstructed in this paper (Fig.2) is favored over the scale-invariant and power-law parametrizations at $\gtrsim 1\sigma$. The results of this paper are completely consistent with the spectrum reconstructed in MW03b using the wavelet band power method. These two methods are very different in nature and complementary (see section 2).

The direct wavelet expansion method has enabled us to find indication of a new localized feature in $P_{in}(k)$ at large scales around $0.001 \lesssim k/\text{Mpc}^{-1} \lesssim 0.005$. It will be interesting to see whether future data will confirm the existence of this feature.

¹¹The χ^2_{eff} of the best fit power-law model for these data is 1043.68 with 6 constrained parameters. In order to compare this model with the wavelet coefficients based model one would need to use a more involved method, possibly that of Bayesian evidence.

We acknowledge helpful discussions with Antony Lewis, Christopher Koenigsberg, and Rich Holman. We thank Henry Neeman for computational support. We acknowledge the use of CAMB and CosmoMC. This work is supported in part by NSF CAREER grant AST-0094335.




Fig. 1.— The figure shows the marginalized 1d distributions of all the parameters used; 16 wavelet coefficients and 3 cosmological parameters. A Gaussian distribution of τ has been marginalized over (see text). The mean values and 1σ standard deviations on the cosmological parameters are $h = 0.59 \pm 0.09$, $\Omega_b h^2 = 0.018 \pm 0.004$ and $\Omega_c h^2 = 0.143 \pm 0.019$. Corresponding constraints on the wavelet coefficients imply constraints on $P_{in}(k)$ shown in Figure 2.




Fig. 2.— reconstructed using all wavelet coefficients (solid curve) with its 1σ confidence region (dotted curves). These constraints are shown overlaid on the constraints obtained by MW03b using the wavelet band power method. (b) $P_{in}(k)$ reconstructed using only those coefficients that deviate from 0 at $> 1\sigma$ (solid curve) with the corresponding 1σ confidence region (dotted curves). These constraints are shown overlaid on the constraints obtained using the tophat binning method with bands carefully positioned closer together so as to be able to pick up the feature at large angular scales around $0.001 \lesssim k/\text{Mpc}^{-1} \lesssim 0.005$, and placed further apart in the remaining parts of the spectrum. The dashed line is the scale-invariant spectrum that best fits these data.

REFERENCES

- Adams, J.A., Ross, G.G., & Sarkar, S. 1997, Nuclear Physics B, 503, 405
- Adams, J., Cresswell, B, & Easter, R. 2001, Phys. Rev. D64, 3514
- Albrecht, A., Steinhardt, P. J. 1982, Phys. Rev. Lett., 48, 1220
- Barger, V., Lee, H.S., & Marfatia, D. 2003, astro-ph/0302150
- Bennett, C., et al. 2003, astro-ph/0302207
- Bridle, S.L., Lewis, A.M., Weller, J., & Efstathiou, G. 2003, astro-ph/0302306
- Bridle, S. L., Crittenden, R., Melchiorri, A., Hobson, M. P., Kneissl, R., & Lasenby, A. N. 2002, MNRAS, 335, 1193
- Chung, D.J.H., Kolb, E.W., Riotto, A., & Tkachev, I.I. 2000, Phys. Rev. D, 62, 043508
- Daubechies, I. 1992, Ten Lectures on Wavelets, S.I.A.M., Philadelphia.
- Efstathiou, G., & Bond, J. R. 1999, MNRAS, 304, 75
- Elgaroy, O.; Gramann, M.; Lahav, O. 2002 MNRAS, 333, 93
- Enqvist, K.; Kurki-Suonio, H. 2000, Phys. Rev. D, 61, 043002
- Feng, B. & Zhang, X. 2003, astro-ph/0305020
- Freese, K., Frieman, J.A., & Olinto, A.V. 1990, Phys. Rev. Lett., 65, 3233
- Gott, J. R. 1982, Nature, 295, 304
- Guth, A.H. 1981, Phys. Rev. D, 23, 347
- Hamilton, A.J.S., Tegmark, M. 2002, MNRAS, 330, 506
- Hannestad, S. 2000, Phys. Rev. D61, 023002
- Holman, R., Kolb, E.W., Vadas, S.L., & Wang, Y. 1991a, Phys. Rev., D43, 3833
- Holman, R., Kolb, E.W., Vadas, S.L., & Wang, Y. 1991b, Phys. Lett., B269, 252
- Hu, W. 2003, Annals of Physics, in press, astro-ph/0210696
- Kinney, W. H. 2001, Phys.Rev. D63, 043001

- Knox, L., Christensen, N., & Skordis, C. 2001, ApJ, 563, L95
- Kogut, A., et al. 2003, astro-ph/0302213
- Kolb, E.W., & Turner, M.S. 1990, *The Early Universe* (Addison-Wesley Publishing Company)
- Kovac, J. et al. 2002, astro-ph/0209478
- Kuo, C.L., et al. 2002, submitted to ApJ, astro-ph/0212289
- La, D., & Steinhardt, P.J. 1991, Phys. Rev. Lett., 62, 376
- Lesgourgues, J., Polarski, D., & Starobinsky, A.A. 1997, Nuclear Physics B, 497, 479
- Lewis, A., Challinor, A., & Lasenby, A. 2000, ApJ, 538, 473, astro-ph/9911177
- Lewis, A., & Bridle, S. 2002, Phys. Rev. D, 66, 103511, astro-ph/0205436
- Linde, A.D. 1983, Phys. Lett., 129B, 177
- Lyth, D. H., Ungarelli, C., & Wands, D. 2002, Phys. Rev. D, in press, astro-ph /0208055
- Mukherjee, P., & Wang, Y. 2003a, ApJ, in press, astro-ph/0301058 (MW03a)
- Mukherjee, P., & Wang, Y. 2003b, astro-ph/0303211
- Neil, R.M. 1993, ftp://ftp.cs.utoronto.ca/pub/radford/review.ps.gz
- Pearson, T.J., et al. 2002, submitted to ApJ, astro-ph/0205388
- Peebles, P.J.E., & Ratra, B. 2003, RMP, in press, astro-ph/0207347
- Peiris, H. V., et al. 2003, astro-ph/0302225
- Percival, W.J. et al. 2002, MNRAS, in press, astro-ph/0206256
- Press, W.H., Teukolsky, S.A., Vetterling, W.T., & Flannery, B.P. 1994, Numerical Recipes, Cambridge University Press, Cambridge.
- Randall, L., Soljagic, M., & Guth, A. 1996, Nucl. Phys. B472, 377
- Rubino-Martin, J.A., et al. 2002, submitted to MNRAS, astro-ph/0205367
- Seljak, U., & Zaldarriaga, M. 1996, ApJ, 469, 437
- Seljak, U., McDonald, P., & Makarov, A. 2003, astro-ph/0302571

- Spergel, D. N.; et al. 2003, astro-ph/0302209
- Tegmark, M., & Zaldarriaga, M. 2002, Phys.Rev. D, 66, 103508
- Verde, L. et al. 2003, astro-ph/0302218
- Wang, Y. 1994, Phys. Rev., D50, 6135
- Wang, Y., Spergel, D.N., & Strauss, M.A. 1999, ApJ, 510, 20
- Wang, Y., & Mathews, G.J. 2001, ApJ, 573, 1